PCT

MIPO PCT

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's	or ag	gent's file reference			N-464-		
11006-98			FOR FURTHER AC			n of Transmittal of International amination Report (Form PCT/IPEA/416)	
International application No.			International filing date (da	ay/month/yea	ır) P	riority date (day/month/year)	
PCT/CA00/00650 02/06/2000				. ,		3/06/1999	
Internation C12N5/0		ent Classification (IPC) or na	tional classification and IPC				
Applicant							
NATION	AL F	RESEARCH COUNCIL	OF CANADA et al.				
 This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36. 							
2. This REPORT consists of a total of 8 sheets, including this cover sheet.							
This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT). These annexes consist of a total of 7 sheets.							
3. This r I II IV V VI VIII		citations and explanation	pinion with regard to nove n der Article 35(2) with reg ns suporting such statem d ternational application	elty, inventiveration and to nove the second		industrial applicability e step or industrial applicability;	
Date of submission of the demand				Date of comp	etion of this r	eport	
12/12/2000				0.09.2001			
Name and mailing address of the international preliminary examining authority:				uthorized of	icer	ad h 60 VES MITEV.	\Box
European Patent Office D-80298 Munich Tel. +49 89 2399 - 0 Tx: 523656 epmu d Fax: +49 89 2399 - 4465				oubradou.	-Bourges,	N (Same Same Same Same Same Same Same Same	K. SUMOTEWN AND THE

INTERNATIONAL PRELIMINARY EXAMINATION REPORT

International application No. PCT/CA00/00650

	an	e receiving Office in d are not annexed to escription, pages:	response to an invitation under to this report since they do not o	r Article 14 are contain amend	referred to in this r Iments (Rules 70.16	eport as "originally filed" 5 and 70.17)):			
	1-3	36	as originally filed						
	Cla	aims, No.:							
	1-5	55	as received on	06/09/2001	with letter of	06/09/2001			
	Dra	awings, sheets:							
	1/1	3-13/13	as originally filed						
2.	Wit lan	h regard to the lang guage in which the i	uage, all the elements marked nternational application was file	above were a ed, unless othe	vailable or furnished erwise indicated und	d to this Authority in the der this item.			
	The	hese elements were available or furnished to this Authority in the following language: , which is:							
		the language of a translation furnished for the purposes of the international search (under Rule 23.1(b)).							
		the language of pu	blication of the international ap	plication (unde	er Rule 48.3(b)).				
		the language of a t 55.2 and/or 55.3).	ranslation furnished for the pur	poses of inter	national preliminary	examination (under Rule			
3.	Witl inte	n regard to any nuc rnational preliminary	leotide and/or amino acid sec y examination was carried out o	quence discloson the basis of	sed in the internation the sequence listin	nal application, the g:			
		contained in the int	ernational application in writter	form.					
		filed together with t	he international application in o	computer read	able form.				
		furnished subsequently to this Authority in written form.							
		furnished subsequently to this Authority in computer readable form.							
		The statement that listing has been fur	the information recorded in connished.	mputer readab	le form is identical t	to the written sequence			
١.	The	amendments have	resulted in the cancellation of:						
		the description,	pages:						
		the claims,	Nos.:						

1. With regard to the elements of the international application (Replacement sheets which have been furnished to



		the drawings,	sheets:					
5.		This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):						
		(Any replacement shoreport.)	eet conta	ining suc	ch amendments must be referred to under item 1 and annexed to this			
6.	Add	litional observations, if	necessa	ıry:				
IV	. Lac	k of unity of inventio	n					
1.	In re	In response to the invitation to restrict or pay additional fees the applicant has:						
		restricted the claims.						
		paid additional fees.						
		paid additional fees u	nder prot	est.				
		neither restricted nor p	paid addi	itional fee	es.			
2.		This Authority found that the requirement of unity of invention is not complied and chose, according to Rule 68.1, not to invite the applicant to restrict or pay additional fees.						
3.	This	s Authority considers that the requirement of unity of invention in accordance with Rules 13.1, 13.2 and 13.3 is						
		complied with.						
		not complied with for t see separate sheet	he follow	ring reaso	ons:			
4.	Con: exan	sequently, the following parts of the international application were the subject of international preliminary nination in establishing this report:						
	☒	all parts.						
		the parts relating to claims Nos						
/ .	Reas citat	soned statement und ions and explanation	er Articl	e 35(2) w rting suc	vith regard to novelty, inventive step or industrial applicability;			
١.	State	ement						
	Nove	elty (N)	Yes: No:		6, 8, 12, 14-31, 34-55 1-5, 7, 9-11, 13, 32-33			
	Inver	ntive step (IS)	Yes: No:	Claims Claims				



International application No. PCT/CA00/00650

Industrial applicability (IA)

Yes:

Claims 1-55

No:

Claims

2. Citations and explanations see separate sheet

VIII. Certain observations on the international application

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made: see separate sheet

Reference is made to the following documents:

- D1: US-A-5 518 915 (NAUGHTON ET AL.) 21 May 1996 (1996-05-21)
- D2: NYGAARD SVEIN J T ET AL: 'Dynamic determination of human glioma invasion in vitro.' JOURNAL OF NEUROSURGERY, vol. 89, no. 3, 1998, pages 441-447, XP000971892 ISSN: 0022-3085
- D3: Steinstrasser I. et al.: "Physical model relating diffusional transport and concurrent metabolism of peptides in metabolically active cell sheets" J. Pharm Sci, 1995 Nov; vol 84 n°11, p. 1332-41
- D4: US-A-5 314 805 (HAUGLAND ET AL.) 24 May 1994 (1994-05-24)

The document D3 was not cited in the international search report. A copy of the document is appended hereto.

SECTION I

The amendments filed with the letter of 06.09.01 are considered to meet the requirements of Art. 34(2) b PCT.

SECTION IV

The present application lacks unity a priori for the following reasons: Claims 1-51 (invention 1) relate to a in vitro model of a mammalian tissue, comprising 3-dimensional aggregates of living mammalian cells. Claims 52-55 (invention 2) relate to a method for predicting biological characteristic of a mammalian tissue.

The IPEA is of the opinion that these two groups of claims are not so linked as to form a single general inventive concept as there is no special technical feature in the sense of Rule 13.2 PCT linking together these two groups of claims, which therefore are considered to be separate inventions

According to Rule 68(1) PCT, the IPEA chose not to invite the applicant to restrict or pay additional fees.

SECTION V

1. **NOVELTY**

The present application does not meet the requirements of Art. 33(2) and (3) PCT:

D2 relates to a study wherein brain tumor tissue is grown in a three-dimensional culture. The tumor cells are cocultured with fetal rat brain cells aggregates. The cells are stained with fluorescent dyes before the coculture.

The cells of the two different phenotypes are considered to be in a "predetermined initial proportion", since equally sized tumor spheroids and brain cell aggregates are used for incubation in the coculture, thus leading to predefined and constant initial proportion of tumor and brain cells (see p.442, Material and Methods, Fluorescent Dye Staining Procedures).

The "proliferation kinetics of the two types of cells" is considered to be "simultaneously assessed" since the tumor proliferation rate and the brain cell aggregates destruction are simultaneously measured and plotted (see p.445, fig.7).

Therefore, D2 is prejudicial to the novelty and the inventive activity of claims 1-5, 7, 9-11, 13, 32-33.

2. INVENTIVE ACTIVITY

Invention 1: The present application relating to invention 1 does not meet the requirements of Art. 33(3) PCT:

2.1 The additional features of dependent claims 6, 8, 12, 14-15, 20-24, 27-30, 34-42 appear to be merely obvious alternatives for a skilled person in the art, and thus are not suitable to establish the presence of an inventive activity in said claims. The teaching of said claims relates to the use of specific fluorescent dyes, the specie and the phenotype of the cells used, etc. All these teachings are anticipated in D1 and D4 for example, which are documents from the same fields, namely three-dimensional culture systems and cell viability assays using dyes.

INTERNATIONAL PRELIMINARY

International application No. PCT/CA00/00650

EXAMINATION REPORT - SEPARATE SHEET

2.2 The use of three-dimensional systems for screening compounds for cytotoxicity, drug action, diagnosing and monitoring cancer is well known and described in the prior art.

For example, D1 (see p. 16 and 32) discloses a three-dimensional model system for the blood-brain barrier comprising tight junctional complexes, which may be used to test the ability of substances to cross the barrier. D1 also states that the three-dimensional systems are useful to study functional and morphological interactions since these systems are mimetic to the physiologic histology and present the cell-cell and cell-matrix interactions. D1 provides complete teaching for such screening.

D2, which is considered to represent the closest prior art to the subject-matter of claims 43-48 and 49-51, gives a clear incentive to use the disclosed threedimensional tissue system as an experimental too for studying the effects of drugs on said system (see p.445, right column) and for evaluating therapy regimens directed against normal brain destruction as well as against biological parameters governing cell migration (see p.446, left column, last §).

Thus, in the light of D2 in combination with the teachings of D1 and the general knowledge of the skilled person in the art, the subject- matter of claims 43-48 and 49-51 which relates to method of screening of antitumour substance and of substance modulating gap junction intercellular interaction does not involve an inventive activity.

Invention 2:

The subject-matter of claims 52-55 appears to be novel and inventive (Art. 33(2) and (3) PCT in view of the available document D3, cited by the examiner for the following reasons:

D3 relates to a method wherein metabolically active cell sheets representing for example the epidermis are provided, parameters like substrate concentration gradients are generated by mathematical simulation, parameters are subjected to variation and thus, metabolism control and diffusion control are predicted. However, D3 fails to provide or suggest a cellular automaton simulation model. Thus, the method of claim 52 and dependent claims thereof is clearly distinguishable and inventive over the mathematical model described in D3.

INTERNATIONAL PRELIMINARY



International application No. PCT/CA00/00650

EXAMINATION REPORT - SEPARATE SHEET

SECTION VIII

The present application does not meet the requirements of Art. 6 PCT:

The term "SIMCEL-2D", "SIMCEL-3D" (claim 53) are internal designation, meaningless to a skilled person in the art.

REPLACEDMO 00/75286 ART 34 AN WHAT IS CLAIMED IS:

5

15



- 1. An *in vitro* model of a mammalian tissue, said model comprising 3-dimensional aggregates of living mammalian cells of at least two different phenotypes and a liquid growth medium.
- 2. A model according to claim 1, wherein the 3-dimensional cell aggregates are suspended in the growth medium.
- 3. A model according to claim 2, wherein the 3-dimensional cell aggregates are of an essentially spheroidal shape.
 - 4. A model according to claim 3, wherein the 3-dimensional cell aggregates are formed in the absence of a solid support.
 - A model according to claim 3, wherein the 3-dimensional cell aggregates are formed in the presence of a solid support.
- 6. A model according to claim 5, wherein the solid support consists of porous beads.
 - 7. A model according to claim 1, wherein the cells of at least one phenotype are fluorescently labeled prior to forming the 3-dimensional aggregates.
- 25 8. A model according to claim 7, wherein the cells are labeled with a fluorescent membrane linker.
 - 9. A model according to claim 7, wherein the cells are labeled by loading with a fluorescent dye.
 - 10. A model according to claim 1, wherein the 3-dimensional aggregates comprise cells of a first and of a second phenotype.



- 11. A model according to claim 10, wherein the cells of at least one phenotype are fluorescently labeled prior to forming the 3-dimensional aggregates.
- 12. A model according to claim 11, wherein the cells are labeled with a fluorescent membrane linker.
 - 13. A model according to claim 11, wherein the cells are labeled by loading with a fluorescent dye.

15

- 14. A model according to claim 13, wherein the dye is calcein-AM.
- 15. A model according to claim 12, wherein the cells of the first and the second phenotype are labeled with fluorescent membrane linkers fluorescing at different wavelengths.
- 16. A model according to claim 10, wherein the cells of the first and the second phenotype are of human origin.
- 17. A model according to claim 16, wherein the cells of the first phenotype are normal cells of human origin.
 - 18. A model according to claim 17, wherein the cells of the first phenotype are endothelial cells.

- 19. A model according to claim 18, wherein the cells of the second phenotype are tumour cells.
- 20. A model according to claim 19, wherein the endothelial cells are fluorescently labeled.
 - 21. A model according to claim 20, wherein the endothelial cells are labeled with a fluorescent membrane linker.



- 22. A model according to claim 21, wherein the 3-dimensional cell aggregates are formed in the absence of a solid support.
- 23. A model according to claim 22, wherein the 3-dimensional cell aggregates are formed by covering particles of a solid support with a layer of the endothelial cells and seeding the tumour organoids to the layer of endothelial cells.
- 24. A model according to claim 23, wherein the solid support is capable of releasing a blood substitute.
 - 25. A model according to claim 17, wherein the cells of the first phenotype are stromal cells.
 - 26. A model according to claim 25, wherein the cells of the second phenotype are tumour cells matching the source of the stromal cells.
- 27. A model according to claim 26, wherein the cells of both phenotypes are fluorescently labeled with labels fluorescing at different wavelengths.
 - 28. A model according to claim 27, wherein the labels are fluorescent membrane linkers.
- 29. A model according to claim 28, wherein 3-dimensional cell aggregates are formed in the absence of a solid support.
 - 30. A model according to claim 28, wherein 3-dimensional aggregates are formed by growing a layer of the stromal cells on particles of a solid support and then growing a layer of the tumour cells on the layer of the stromal cells.

10



- 31. A model according to claim 17, wherein the cells of the second phenotype are tumour cells.
- 32. A model according to claim 13, wherein the cells of the first phenotype are cells of a tissue in which metastases of the tumour are expected to develop.
- 33. A model according to claim 32, wherein the cells of the first phenotype are epithelial cells.
- 34. A model according to claim 31, wherein the cells of the first phenotype are grown as a monolayer on one side of a porous solid support.
- 35. A model according to claim 34, wherein the tumour cells grown in the form of 3-dimensional aggregates are applied to the opposite side of the support.
 - 36. A model according to claim 35, wherein the tumour cells are fluoroscently labeled.
 - 37. A model according to claim 17, wherein the cells of the second phenotype are cells of the first phenotype treated with a chemical agent prior to forming the 3-dimensional aggregates.
- 25 38. A model according to claim 37, wherein the chemical agent is capable of blocking the proliferation of cells without killing the cells.
 - 39. A model according to claim 38, wherein the chemical agent is mitomycin.
- 30 40. A model according to claim 37, wherein the chemical agent is a phototoxic agent.

10

15

20



- 41.A model according to claim 40, wherein the chemical agent is chloromethyl eosine diacetate.
- 42. A model according to claim 40, wherein the 3-dimensional aggregates of cells are illuminated with a light source after formation.
 - 43. A method of screening for an antitumour substance, said method comprising the steps of:
 - a. providing a 3-dimentional *in vitro* model of human tissue according to claim 1, said model comprising at least one phenotype of tumour cells:
 - b. providing a candidate antitumour substance;
 - c. culturing the 3-dimensional cell aggregates for a predetermined period of time, in the presence and in the absence of the candidate antitumour substance;
- d. measuring the cell proliferation rate of at least one cell phenotype in the 3-dimensional cell aggregates in the absence and in the presence of the candidate antitumour substance; and
 - e. accepting or rejecting the candidate antitumour substance based on results of the measurements of step d.
- 44. A method according to claim 43, wherein cells of at least one cell phenotype are fluorescently labeled.
- 45. A method according to claim 44, wherein cells are labeled with a fluorescent membrane linker.
 - 46. A method according to claim 45, further including the step of dispersing the cell aggregates into a suspension of individual cells prior to measuring the cell proliferation rate.
 - 47. A method according to claim 46, wherein the proliferation rate is expressed as the proliferation index.

20



- 41.A model according to claim 40, wherein the chemical agent is chloromethyl eosine diacetate.
- 42. A model according to claim 40, wherein the 3-dimensional aggregates of cells are illuminated with a light source after formation.
 - 43. A method of screening for an antitumour substance, said method comprising the steps of:
 - a. providing a 3-dimentional *in vitro* model of human tissue according to claim 1, said model comprising at least one phenotype of tumour cells;
 - b. providing a candidate antitumour substance;
 - c. culturing the 3-dimensional cell aggregates for a predetermined period of time, in the presence and in the absence of the candidate antitumour substance;
- d. measuring the cell proliferation rate of at least one cell phenotype in the 3-dimensional cell aggregates in the absence and in the presence of the candidate antitumour substance; and
 - e. accepting or rejecting the candidate antitumour substance based on results of the measurements of step d.
 - 44. A method according to claim 43, wherein cells of at least one cell phenotype are fluorescently labeled.
- 45. A method according to claim 44, wherein cells are labeled with a fluorescent membrane linker.
 - 46. A method according to claim 45, further including the step of dispersing the cell aggregates into a suspension of individual cells prior to measuring the cell proliferation rate.
 - 47. A method according to claim 46, wherein the proliferation rate is expressed as the proliferation index.

- 48. A method according to claim 47, wherein the proliferation index is calculated from flow cytometry analysis of the cell suspension.
- 49. A method of screening for a substance modulating gap junction intercellular communication, said method comprising the steps of:
 - a. providing a 3-dimensional in vitro model of human tissue according to claim 1, said model comprising at least cell phenotype loaded with a fluorescent dye impermeant to the cell membrane;
- b. providing a candidate modulating substance;
 - c. culturing the 3-dimensional cell aggregates for a predetermined period of time, in the presence and in the absence of the candidate substance:
 - d. measuring the migration of the dye to at least one other cell phenotype in the 3-dimensional cell aggregates, in the absence and in the presence of the candidate antitumour substance; and
 - e. accepting or rejecting the candidate modulating substance based on results of the measurements of step d.
- 50. A method according to claim 49, further including the step of dispersing the cell aggregates into a suspension of individual cells prior to measuring the migration of the dye.
 - 51. A method according to claim 50, wherein the dye is calcein-AM.
 - 52. A method for predicting a biological characteristic of a mammalian tissue, said method comprising the steps of:
 - a. providing a simulation model of the mammalian tissue;
 - b. setting model parameters;
- 30 c. running the model; and
 - d. evaluating the biological characteristic of the tissue based on results of step c.

- 53. A method according to claim 51, wherein the simulation model is SIMCEL-2D or SIMCEL-3D simulation model.
- 54. A method according to claim 52, wherein the biological response is the cyclic cell recruitment from the resting pool, the cell proliferation index, or intercellular diffusion.
- 55. A method according to claim 53, wherein the model parameters are set to simulate effects of physical or chemical agents on the tissue.